

## **SEMINÁRIO DE ANÁLISE E EQUAÇÕES DIFERENCIAIS**

**Dia 12 de Julho (quinta-feira), às 13H30, na sala 6.2.33**

### **The Cheeger problem and an application to the (constant) Prescribed Mean Curvature problem**

**Giorgio Saracco**

(Università degli Studi di Pavia)

#### **Abstract**

Given an open, bounded set  $\Omega$ , one defines its Cheeger constant  $h(\Omega)$  as the infimum of the ratio perimeter over volume among all of its subsets. Evaluating  $h(\Omega)$  and finding the sets  $E$  that attain such a minimum is known as the Cheeger problem.

There are many possible motivations to study such a problem as the constant  $h(\Omega)$  and minimizers of the ratio play a major role in different areas. In particular we shall discuss the connection with the (constant) prescribed mean curvature problem giving a characterization of existence and uniqueness of solutions in terms of the Cheeger problem.

It will be clear that being able to compute  $h(\Omega)$  and knowing who the minimizers are is of interest. In general though these are difficult tasks, even in the planar case. We shall show that for a class of a Jordan domains there is a structure theorem for minimizers. On top of that, the so-called inner Cheeger formula holds and this allows to compute the exact value of  $h(\Omega)$ .

These results have been obtained in collaboration with G.P. Leonardi (Università di Modena e Reggio-Emilia, IT) and R. Neumayer (Northwestern University, USA)

*Seminário financiado por Fundos Nacionais através da FCT – Fundação para a Ciência e a Tecnologia no âmbito do projeto UID/MAT/04561/2013*